

Revealing the Large-Scale Structures of Interstellar Gas Associated with the Magellanic SNR N132D

H. Sano¹, Y. Fukui¹, S. Yoshiike¹, T. Fukuda¹, K. Tachihara¹, S. Inutsuka¹, A. Kawamura², K. Fujii², N. Mizuno², T. Inoue², T. Onishi³, F. Acero⁴, J. Vink⁵

¹*Department of Physics, Nagoya University, Furo-cho, Chikusa-ku, Nagoya 464-8601, Japan; sano@a.phys.nagoya-u.ac.jp*

²*National Astronomical Observatory of Japan, Mitaka 181-8588, Japan*

³*Department of Physical Science, Graduate School of Science, Osaka Prefecture University, 1-1 Gakuen-cho, Naka-ku, Sakai 599-8531, Japan*

⁴*Laboratoire AIM, CEA-IRFU/CNRS/Université Paris Diderot, Service d'Astrophysique, CEA Saclay, 91191 Gif sur Yvette, France*

⁵*Astronomical Institute Anton Pannekoek, University of Amsterdam, P.O. Box 94249, 1090 GE Amsterdam, The Netherlands*

Abstract. We report preliminary results of large-scale distribution toward the Magellanic supernova remnant N132D using Mopra and *Chandra* archival datasets. We identified a cavity-like CO structure along the X-ray shell toward the southern half of it. The total mass of associating molecular gas is $\sim 10^4 M_\odot$, which is smaller than the previous study by an order of magnitude. Further observations using ALMA, ASTE, and Mopra will reveal the detailed spatial structures and its physical conditions.

1. Introduction

In supernova remnants (SNRs), interaction between shock waves and surrounding interstellar gas is a key element for understanding the SNR evolution, cosmic ray acceleration, and origin of gamma-ray radiation. Recently, Sano et al. (2015) revealed that the shock-cloud interaction in a young Galactic SNR generates the turbulence and strong magnetic field, which enhance the non-thermal X-rays and an efficient acceleration of the cosmic ray electrons around the interacting gas clumps. Therefore, it is important to search for more evidence to examine its universality.

The SNR N132D is a bright X-ray emitter in the Large Magellanic Cloud (LMC; Figure 1a), and identified as a young SNR (~ 3150 yr). It is also known as an Oxygen-rich SNR (e.g., Borkowski et al. 2007) and is hence considered to be a remnant of a core-collapse supernova of a massive star. Furthermore, the SNR is thought to be possibly interacting with molecular gas as shown by CO observations with SEST (Banas et al. 1997). Therefore, it is the best target to study the shock-cloud interaction in the extra galaxies. In the present paper, we report the large-scale distribution of the associate molecular gas with N132D and its physical properties.

2. Results

Figure 1a shows *Chandra* X-ray tricolor image of N132D (Borkowski et al. 2007), which has shell-like and filamentary structures especially in the southwest, while the blowout structure appears in the northeast. Thermal X-rays (corresponding to a energy




Figure 1. (a) The *Chandra* X-ray tricolor image of SNR N132D (Borkowski et al. 2007). (b) Integrated intensity map of $^{12}\text{CO}(J=1-0)$ (Wong et al. 2011) in a velocity range of $V_{\text{LSR}} = 256.5\text{--}268.6 \text{ km s}^{-1}$ is shown in color (unit is K km s^{-1}). The white contours correspond to the *Chandra* X-ray flux in the energy band 0.5–7.0 keV.

range of 0.5–1.2 keV) are bright over the whole SNR, whereas non-thermal X-rays (2.0–7.0 keV) emit only in the southern half.

Figure 1b represents the $^{12}\text{CO}(J=1-0)$ integrated intensity map taken by the Magellanic Mopra Assessment (MAGMA; Wong et al. 2011) with the Mopra 22-m telescope. The existence of the giant molecular cloud (GMC) in the southern part of Figure 1b was known by a previous study (Banas et al. 1997). We newly found three molecular clouds interacting with the SNR. The two clouds are located in the west and the east, which form the cavity-like CO structure along the X-ray shell. Another CO cloud is located in the northeast. The total molecular mass of the newly found 3 clouds is $\sim 10^4 M_{\odot}$ using the X -factor $7 \times 10^{20} [W(^{12}\text{CO})/(\text{K km s}^{-1})] (\text{cm}^{-2})$ (Fukui et al. 2008), which is smaller than the previous study by an order of magnitude (Banas et al. 1997). It is because the most part of the GMC is not interacting with the SNR.

3. Discussion and Summary

We discussed that the CO cavity-like structure was created by the stellar wind / UV photons from the massive star prior to the supernova explosion and is now interacting with the SNR shocks. Furthermore, the enhancement of non-thermal X-rays in the southern part can be described as a result of shock interaction with clumpy CO structures (e.g., Sano et al. 2015). Therefore, we predict that the surroundings of N132D have clumpy CO structures corresponding to the non-thermal X-ray filaments. We continue follow up observations with ALMA, ASTE, and Mopra, which allow us for the first time to study the interaction of a SNR with CO gas in detail outside our own Galaxy.

References

- Sano, H., Fukuda, T., Yoshiike, S., et al. 2015, *ApJ*, 799, 175
- Borkowski, K. J., Hendrick, S. P., & Reynolds, S. P. 2007, *ApJL*, 671, L45
- Banas, K. R., Hughes, J. P., Bronfman, L., & Nyman, L.-Å. 1997, *ApJ*, 480, 607
- Wong, T., Hughes, A., Ott, J., et al. 2011, *ApJS*, 197, 16
- Fukui, Y., Kawamura, A., Minamidani, T., et al. 2008, *ApJS*, 178, 56